



# Comorbidity Implications in Brain Disease: Neuronal Substrates of Symptom Profiles

TOMAS PALOMO<sup>a</sup>, RICHARD J. BENINGER<sup>b</sup>, RICHARD M. KOSTRZEWA<sup>c</sup> and TREVOR ARCHER<sup>d,\*</sup>

<sup>a</sup>Psychiatry Service, "12 de Octubre", University Hospital, Madrid 28041, Spain; <sup>b</sup>Departments of Psychology and Psychiatry, Queen's University, Kingston ON, K7L 3N6 Canada; <sup>c</sup>Department of Pharmacology, Quillen College of Medicine, East Tennessee State University, Johnson City, TN USA; <sup>d</sup>University of Göteborg, Department of Psychology, Box 500, SE-45030 Göteborg, and University of Kalmar, HBV, SE-39182 Kalmar, Sweden. [trevor.archer@psy.gu.se](mailto:trevor.archer@psy.gu.se)

(Submitted 19 January 2006; In final form 19 January 2006)

**The neuronal substrates underlying aspects of comorbidity in brain disease states may be described over psychiatric and neurologic conditions that include affective disorders, cognitive disorders, schizophrenia, obsessive-compulsive disorder, substance abuse disorders as well as the neurodegenerative disorders. Regional and circuitry analyses of biogenic amine systems that are implicated in neural and behavioural pathologies are elucidated using neuroimaging, electrophysiological, neurochemical, neuropharmacological and neurobehavioural methods that present demonstrations of the neuropathological phenomena, such as behavioural sensitisation, cognitive impairments, maladaptive reactions to environmental stress and serious motor deficits. Considerations of neuronal alterations that may or may not be associated with behavioural abnormalities examine differentially the implications of discrete areas within brains that have been assigned functional significance; in the case of the frontal lobes, differential deficits of ventromedial and dorsolateral prefrontal cortex may be associated with different aspects of cognition, affect, remission or response to medication thereby imparting a varying aspect to any investigation of comorbidity.**

**Keywords:** Neuronal systems; Regional selectivity; Cognition; Affect; Neuroimaging; Signalling; Sensitisation; Stress; Comorbidity

## INTRODUCTION

Comorbidity implications within and between the neurologic and neuropsychiatric disorders remain, and seem even aggravated, by calamities of high incidence with consequential problems for both the etiology and treatment of these states (Walker and Diforio, 1997). Commonly, patients afflicted with psychiatric disorders, such as depression or schizophrenia, are at high risk for any of several substance abuse disorders, including alcohol, tobacco or unlawful substances (Colder and Chassin, 1993; Grant BF, 1995; George and Krystal, 2000; Chambers *et al.*, 2001; Dierker *et al.*, 2002; Ziedonis *et al.*, 2003), while affective and psychotic disorders may present comorbidity with disorders of cognition (Golik-Gruber *et al.*, 2003; Lewis *et al.*, 2004). In view of the occurrence of particular shared genetic liabilities for comorbidity (Kendler *et al.*, 1993; Hambrecht and Hafner, 2000; Johnson JG *et al.*, 2000; Swendsen and Merikangas, 2000; Katarina *et al.*, 2004), the pres-

\*Corresponding author: E-mail: [trevor.archer@psy.gu.se](mailto:trevor.archer@psy.gu.se)

ence of shared neuromorphological and anatomical features of these disorders (*e.g.*, Akbarian *et al.*, 1993; Pierri *et al.*, 1999; Drevets, 2000; Glantz and Lewis, 2000; Eastwood *et al.*, 2001). Further, the shared features of affective, psychotic and neurologic states encompass a variety of neurobiological markers (Buchanan *et al.*, 1998; Woo *et al.*, 1998; Rajkowska *et al.*, 1999; Harrison *et al.*, 2003; Coryell *et al.*, 2005), as well as the central role of neurodevelopmental prerequisites from a genetic-environmental perspective (Weinberger, 1987; Stine *et al.*, 1993; Campodonico *et al.*, 1998; Feigin *et al.*, 2001; Chambers *et al.*, 2003), all of which underline both the sufficiency and necessity of a consideration of the neuronal substrates of comorbidity.

### NEUROTRANSMITTER-REGIONAL SELECTIVITY

The biogenic amines would appear to be implicated in several aspects of brain comorbid pathology with dopaminergic and serotonergic systems involved in schizophrenia, the serotonergic and noradrenergic systems in depression and anxiety disorders, dopaminergic influences in anhedonia and differential affectivity, and the maintenance of substance abuse and drug dependence (Eastwood and Harrison, 2001; Davidson *et al.*, 2002; Smelson *et al.*, 2002; Kroner *et al.*, 2005). Furthermore, the aminergic contributions have been associated with several anatomically and functionally predominant regions of the brain, including the frontal lobe, above all the prefrontal and orbitofrontal areas, hippocampus, amygdala, anterior cingulate cortex, striatum, thalamus and nucleus accumbens (O'Donnell and Grace, 1995; Mayberg, 1997; Gray, 1999; Groenewegen *et al.*, 1999; LeDoux, 2000), and the prerequisite of neuronal plasticity (Spedding *et al.*, 2003; Kauer, 2004). Totterdell outlined anatomically converging dopaminergic and cortical inputs to single neurons located in the ventral striatum, as well as the convergence between different cortical inputs onto individual neurons (*cf.*, Wright and Groenewegen, 1995; Totterdell and Meredith, 1997; Rosenkranz and Grace, 2001). Thus, there is a 'gating' mechanism with excitatory inputs from the hippocampus and prefrontal cortex to the ventral element of the prefrontal-striatal-pallidal-thalamic-prefrontal

loop, with major dopaminergic inputs from the ventral tegmentum and serotonergic innervation from the raphe nuclei. Against this regional-neuronal template, descriptions of the synaptic circuitry and excitatory inputs to the nucleus accumbens (Aylward and Totterdell, 1993; Ding *et al.*, 2001), its dopaminergic, serotonergic and noradrenergic innervation and the aminergic receptor distributions were provided. Afferent inputs from the hippocampus, prefrontal cortex, and amygdala (Totterdell and Smith, 1989; Callaway *et al.*, 1991; Johnson LR *et al.*, 1994; Wright *et al.*, 1996; Carr *et al.*, 1999; French and Totterdell, 2002; 2003) maintain the projections underlying the synthesis of pathways involved in drug dependence disorders, depressive disorders, and schizophrenia (*e.g.*, Totterdell and Smith, 1986; Sapolsky, 2001; Crombag *et al.*, 2005; Totterdell, 2006).

High level structural and functional resolution offered by functional magnetic resonance imaging (fMRI), markedly superior to other neuroimaging techniques, provides direct estimates of neuronal activity, thereby information on neuronal substrates, pertaining to aspects of comorbid pathology in brain disorders (*cf.*, Aber *et al.*, 2003; McClure *et al.*, 2004). For example, structural and functional imaging studies, described by Ebmeier *et al.* (2006), implicate the anterior cingulate cortex, orbitofrontal cortex, dorsolateral frontal cortex, striatum and medial temporal areas in abnormalities associated with depressive states (Ebert and Ebmeier, 1996). Thus, depressed patients presented abnormally increased activity in emotion-associated region and abnormal hypoactivity in cognition-associated regions (Drevets, 2000; Ebmeier and Kronhaus, 2002; Mayberg, 2003; Steele and Lawrie, 2004). It was found that spatial working memory performance was linked to abnormal increments in ventromedial prefrontal activity (Rose *et al.*, 2005), consistent with an 'emotional gating' model of cognitive performance in depressive states (Pochon *et al.*, 2002). It was shown too that depressed patients displayed a bias towards processing mood-congruent information, *i.e.*, an elevated response to stimuli with negative emotional tone, whereby the responses to "sad targets" in the rostral anterior cingulate and medial prefrontal cortex were elevated (Elliott *et al.*, 2002). Other studies demonstrated that patients engaged in prolonged, elaborative

cognitive processing of emotional material (Siegler *et al.*, 2002). Taken together, a growing consensus supports the convergence of an affective-cognition component of symptom profile, not least executive functioning (Harvey *et al.*, 2004), as reflected by neuroimaging and neuropsychological studies (Shah *et al.*, 1998; Elliott *et al.*, 2000; Austin *et al.*, 2001; Fossati *et al.*, 2002; 2004; Murphy *et al.*, 2003; Porter *et al.*, 2003; Steele *et al.*, 2004).

The electrophysiological basis of potent prefrontal cortex (PFC) regulatory influences upon limbic regions, particularly the amygdala, in affective and psychotic conditions and the effects of stress were described by Grace (2006). The PFC regulation of subcortical systems involved in affective states has received much attention (Gresch *et al.*, 1994; Jedema *et al.*, 1999; Quirk *et al.*, 2000; Milad and Quirk, 2002; Correll *et al.*, 2005; Goto and Grace, 2005; 2006; Laviolette *et al.*, 2005; Valenti and Grace, 2005). Besides a functioning PFC-amygdala circuit regulation in affective states (Hariri *et al.*, 2003; Rosenkranz and Grace, 2003), hippocampus-PFC interactions within the nucleus accumbens are critical such that induction of long-term potentiation (LTP) interferes with the afferent drive to the accumbens from the other pathway (*i.e.*, PFC or hippocampus) with dopamine exerting modulatory control (Rosenkranz and Grace, 2002a,b). For example, in cocaine-sensitized animals, LTP is induced in the hippocampal-accumbens pathway with consequent disruption of the PFC-accumbens pathway control over goal-directed behaviour (*cf.*, Grace *et al.*, 1998). Genetic-environmental factors contributing to comorbidity, *e.g.*, chronic/acute stress, affect normally-functioning and dysfunctional brain at several different levels and regional sites, most particularly neuronal (Rohsenow, 1982; Abercrombie and Jacobs, 1987; Dunn, 1988; Coco *et al.*, 1992; Romelsjo and Hasin, 1992; Finlay *et al.*, 1995; 1997; Sapolsky, 1996; Mana and Grace, 1997; Caspi *et al.*, 2003; Higihara *et al.*, 2003; Gilder *et al.*, 2004). For instance, exposure to long-term stress induced a remarkable and long-lasting inhibition of LTP that could be prevented by the glucocorticoid receptor antagonist, mifepristone, thereby demonstrating the essential role of not only the hippocampus (as described by Sapolsky, 2005; but see also Sapolsky *et al.*, 1990; Nisenbaum *et al.*, 1991; Jedema *et al.*, 2001) but also the frontal

cortex in the stress response cascade, as well as dopaminergic activity in the ventral tegmentum (Moore *et al.*, 2001). Additionally, both antidepressant compounds, *e.g.*, fluoxetine and tianeptine, and clozapine, but not haloperidol, restored stress-disrupted LTP, suggesting that the effective compounds share the PFC as their common target by restoring the functional balance at hippocampus/PFC synapses, presumably impaired in depression and/or schizophrenia.

Szechtman and co-workers (*e.g.*, Szechtman and Woody, 2004; 2006; Taylor *et al.*, 2005; Woody and Szechtman, 2005; Woody *et al.*, 2005), taking as their point-of-departure the remarkably high prevalence of comorbidity of obsessive-compulsive disorder in patients presenting other psychiatric problems (*e.g.*, Tukul *et al.*, 2002), have developed the notion that the behavioural abnormalities expressed in the disorder (*i.e.*, obsessive-compulsiveness) stem from disturbances in the operation of a "security-motivation" system. Incorporated within the "security-motivation system", and encompassing presumably orbitofrontal-cingulate-limbic-motor elements of regional circuitry, exists a representative, biologically-primitive 'specialized' motivation, activated by potential (rather than imminent) danger signals to 'self' or 'intimate-others', that engages a set of specialized species-typical behaviours that are meant to handle the threat. To attain a "switching-off" of the security motivation system, a self-generated affective state of knowledge or particular insight must occur, a satiety signal termed "Yedasentience". The implications of affective 'constitution' appear to be of central influence for the notions that have been developed to a global view of a 'para-disordered' brain wherein the search for the neuronal substrates is under mobilization (Davidson *et al.*, 2000); implications of affect for basic aspects of personality (*e.g.*, affective, state-dependent, personality, or trait-dependent personality) must be tackled (Palomo *et al.*, 2004). Obsessive-compulsive disorder develops from failure to generate or respond to the yedasentience signal: thus, patients persist in the prolonged repetitive sequences of security-related behaviours (checking, washing, locking, unlocking, counting) reflecting the strong motivational state characteristic of the disorder. The authors have drawn attention also to the concept of "affective chronometry", that mea-

sure the temporal dynamics of emotions/affect within parameters of threshold, peak amplitude and recovery time, that may be applied to assess malfunctioning within the "security-motivation" system or "Yedasentience". From the viewpoint of notions suggesting that obsessive-compulsive disorder may precede the diagnosis of other psychiatric disease states such as schizophrenia as well as with a predisposing incidence of Tourette's disorder, and other movement-oriented disorders, one may imply that the same gene is responsible for tics and obsessive-compulsive disorder; these various threads must reinforce implications of the relations to obsessional repetitive, or more or less stereotypic behaviours (*cf.*, Bolton *et al.*, 2000), all of which serves to impart a Braakian perspective (*cf.*, Modell *et al.*, 1989).

The comorbidity of obsessive-compulsive disorder with other psychiatry disorders is illustrated persuasively in its shared comorbidity with Trichotillomania (TTM), characterized by repetitive stereotypical hair-pulling from different sites - resulting in noticeable hair loss (Stein *et al.*, 1995; Tukul *et al.*, 2001). Both TTM and obsessive-compulsive disorder patients describe compulsive urges and ritualistic behaviours, yet marked differences exist between the disorders, *e.g.*, hair-pulling in TTM is not in response to obsessive thoughts (such as worry about harm to self or others) but rather because of an irresistible urge and the promise of gratification when pulling out hair. Furthermore, TTM maintains the hair-pulling without evolving non-self-injurious compulsive rituals, symptoms that develop chronically with regard to focus and severity, *e.g.*, from washing of hands to checking locks, stoves, appliances, etc. (Swedo and Leonard, 1992). Recently, Lochner *et al.* (2005) compared 278 male and female patients presenting obsessive-compulsive disorder with 54 patients presenting TTM over all age groups. They found that whereas obsessive-compulsive disorder patients presented significantly greater disability over lifetime, TTM patients reported a lesser response to treatment. Obsessive-compulsive disorder patients presented a higher level of comorbidity, greater harm avoidance and less novelty seeking, a higher number of maladaptive beliefs, and more sexual abuse than TTM patients. The study by Lochner *et al.* (2005) serves to underline problematic aspects of comorbidity: TTM

is classified as an impulse control disorder and may resemble conditions characterized by stereotypical self-injurious symptoms, such as skin-picking, thereby implying that important differences may exist in the underlying neuronal-regional circuitry which necessitate contrasting treatment approaches (Keuthen *et al.*, 1998; O'Sullivan *et al.*, 2000; van Minnen *et al.*, 2003).

## NEUROCHEMICAL-REGIONAL SELECTIVITY IN COGNITION AND MOVEMENT

Comorbidity of functional status and regional-neuronal integrity implicates hippocampal, amygdaloid, medial septal and cortical, medial prefrontal and entorhinal involvement and participation in aversive memories through mechanisms implying glutamate receptor activation both during the consolidation phase and during the expression of memory (*e.g.*, Bianchin *et al.*, 1994). The cascade of molecular events underlying retrieval of a simple step-down conditioning task and the memory for resistance to extinction involving functional NMDA receptors in addition to activation of several signalling pathways in the hippocampus, including those mediated by PKA, ERK1/2, p38MAPK and Src provides markedly stringent analysis of neuronal molecular substates of fear and avoidance memory acquisition and extinction (Bernabeu *et al.*, 1995; Izquierdo and Medina, 1995; Maren and Quirk, 2004). Ivan Izquierdo *et al.* (2006) compared problems of memory arising from temporo-limbic and striatal malfunctioning by regarding instances of declarative or procedural memory tasks. His two major investigative thrusts involved: (i) reversal learning in the circular water maze task whereby the declarative component is altered but the procedural one maintained (but see also reference versus working memory), and (ii) inhibitory avoidance conditioning studies suggesting the 'shift of memories' over distributed systems due to tasks initiating further learning. Collectively, these studies revealed how different brain structures and signalling pathways contributed to multiple cognitive functions (*e.g.*, Medina *et al.*, 1999). Further evidence for the neurochemical cascades underlying memories for fear conditioning is offered by a series of experiments reviewed by Cammarota *et al.* (2007), whereby

activity-dependent alterations in neuronal efficacy are considered the main events in the formations and storage of new memories (Dudai, 2002; Izquierdo and Medina, 1997; McGaugh, 2000). They demonstrated that the 'superfamily' of mitogen-activated protein kinases (MAPKs) contribute to signal transduction cascades initiated by neurotransmitters, mitogens, growth factors and various types of stress (*e.g.*, Alonso *et al.*, 2003; McGaugh and Roozendaal, 2002). It was demonstrated that modifications of the phosphorylation/activation state of different synaptic and nuclear-localized protein kinases influenced, by a variety of mechanisms, both the induction and maintenance of neuronal expressions of plasticity, as well as the cellular mechanisms underlying consolidatory processes in the acquisition of long-term memories (*cf.*, Berman DE and Dudai, 2001; Dudai, 2004). Rodriguez-Moreno *et al.* (2006) has applied electrophysiological recording techniques during acquisition and retrieval of cognitive tasks to study associative learning in mice. Activity-dependent changes occur at the hippocampal CA3-CA1 synapse in behaving mice during the acquisition and extinction of a trace conditioning paradigm. A single pulse presented at the Schaffer collateral-commisural pathway, during the CS-US interval, evoked a monosynaptic field EPSP at ipsilateral CA1 pyramidal cells, thereby altering the slope across learning. LTP evoked by train stimulation of Schaffer collaterals blocked acquisition of the associative task. Applications of NMDA antagonists were critical in preventing LTP induction, acquisition of learned eyelid responding and synaptic changes at CA3-CA1 synapses across conditioning. These experimental paradigms were tested also in transgenic (APP, PS-1) and knock-out (TrkB) mice.

Fredriksson and Archer (2006) described the influence of postnatal iron overload upon implications of the functional and interactive role of dopaminergic and noradrenergic pathways that contribute to the expressions of movement disorder. Postnatal iron overload at doses of 7.5 mg/kg (administered on Days 10-12 post partum) and above, invariably induced a behavioural syndrome consisting of an initial (1st 20-40 min of a 60-min test session) hypoactivity followed by a later (final 20 min of a 60-min test session) hyperactivity, when the mice were tested at adult ages (age 60 days or more).

Following postnatal iron overload, subchronic treatment with the neuroleptic compounds, clozapine and haloperidol, dose-dependently reversed the initial hypoactivity and later hyperactivity induced by the metal. Furthermore, DA D<sub>2</sub> receptor supersensitivity (as assessed using the apomorphine-induced behaviour test) was directly and positively correlated with iron concentrations in the basal ganglia. The selective denervation of noradrenaline (NA) terminals using the selective NA neurotoxin, DSP4, followed by administration of the selective DA neurotoxin, MPTP, has been employed as an experimental model for 'accelerated' Parkinson's disease (PD), reflecting a striatal DA and central NA deficiency (Nishi *et al.*, 1991; Marien *et al.*, 1993; Fornai *et al.*, 1997). Brain noradrenaline (NA) denervation, using the selective NA neurotoxin, DSP4, prior to administration of the selective DA neurotoxin, MPTP, exacerbated both the functional (hypokinesia) and neurochemical (DA depletion) effects of the latter neurotoxin. Treatment with L-Dopa restored motor activity only in the animals that had not undergone NA denervation. Finally, C57/BL6 mice were administered either postnatal iron (Fe<sup>2+</sup> 7.5 mg/kg, on postnatal days 10-12) or vehicle, followed by administration of either DSP4 (50 mg/kg, *s.c.*, 30 min after injection of zimeldine, 20 mg/kg, *s.c.*) or vehicle (saline) at 63 days of age. Postnatal iron administration exacerbated the bradykinesia induced by MPTP and virtually abolished all spontaneous motor activity in NA-denervated mice that were MPTP-treated. Suprathreshold doses (20 mg/kg) of L-Dopa invariably restore motor activity in MPTP mice: nevertheless, postnatal iron administration reduced markedly the restoration of motor activity by suprathreshold L-Dopa (20 mg/kg) following a 60-min habituation to the test chambers. Pretreatment with DSP4 effectively eliminated the restorative effect of L-Dopa in the MPTP mice. Postnatal iron administration caused enduring higher levels of total iron content in all the groups with an increased level in mice treated with DSP4 followed by MPTP. From a perspective of Braakian notions, these divergent findings confirm the developmental and predispositional role of postnatal iron overload and prior denervation of NA upon dopaminergic functional expression and indicate a permanent vulnerability both in the noradrenergic and dopaminergic pathways following

the postnatal infliction of an iron overload.

### IMMUNOLOGICAL-REGIONAL FUNCTIONING

The review by Müller and Schwarz (2006) presented a series of bridges over current notions of psychoneuroimmunological functioning and diverse findings from pharmacological, neurochemical and genetic studies implicating glutamatergic-dopaminergic substrates of comorbidity in schizophrenia (Carlsson *et al.*, 2001; Collier and Li, 2003; Kornhuber *et al.*, 2004). One aspect of dysfunctional states concerned the role of glutamatergic hypofunctioning, as mediated by NMDA-R antagonism. Other sources of etiological predisposition pertained to perinatal infection in the pathogenesis of schizophrenia (Hill, 1999; Cook and Hill, 2001; Müller, 2004) and inflammation (Leonard and Myint, 2006). The incidence of pregnancy-infections associated with later susceptibility for has been established (Mednick *et al.*, 1988; Westergaard *et al.*, 1999; Buka *et al.*, 2001; Brown *et al.*, 2004; Koponen *et al.*, 2004). Early sensitization within the immune system may become expressed by an imbalance between Type I and Type II immune responses: the former partially inhibited and the latter over-activated (Müller *et al.*, 2000; Schwarz *et al.*, 2001). Inhibition of indoleamine dioxygenase, an enzyme located in astrocytes and microglia, is inhibited by Type II cytokines so that tryptophan is predominantly metabolised by tryptophan 2,3-dioxygenase in the astrocytes that lack the neurochemical agents for normal tryptophan metabolism. Analysis of the frontal cortex tissue of schizophrenic patients indicated an elevated expression of tryptophan 2,3-dioxygenase in comparison with indoleamine dioxygenase, which was evidenced in the astrocytes, rather than microglial cells (Miller *et al.*, 2004), thereby implying the involvement of glial entities in the Type I/Type 2 imbalance (Aloise *et al.*, 1997; Xiao and Link, 1999). Astrocytes, microglial cells and macrophages were implicated in the cellular localization of kynureneine metabolism (Espesy *et al.*, 1997; Speciale and Schwarcz, 1993). The genetic variations and multiple functions of neuregulin-1 in a diversity of neuronal and glial tissues were implicated too in the etiology of disease process in schizophrenia (Stefansson *et al.*,

2002; 2003; Schwab *et al.*, 2003; Williams *et al.*, 2003; Numakawa *et al.*, 2004). Lack of kynurate-OHase in astrocytes allows build-up of kynurenate acid (see above, endogenous CNS NMDA-R antagonist) which is found together with an increase in tryptophan 2,3-dioxygenase activity in the CNS of schizophrenics (*cf.*, Schwarcz *et al.*, 2001). Finally, the treatment profiles of neuroleptic compounds upon cellular immune reactivity in schizophrenia was discussed (Müller *et al.*, 1991; 1997a,b; 1999; 2002). It would appear then that the immune-mediated glutamatergic-dopaminergic imbalance may be circumvented through selective intervention by anti-inflammatory cyclo-oxygenase-2 inhibitors that provide direct reduction of kynurenate acid to ameliorate the disease process.

Abnormal frontal cortical functioning (Tollefson, 1996; Gehring and Willoughby, 2002), including attention, memory and executive functioning, is observed in a wide range of neuropsychological impairments, such as set-shifting, concentration, judgement, planning and anticipation, abstract thinking, selective attention, mental sequencing and concept formation in schizophrenic individuals (Goldberg *et al.*, 1988, 1990, 1993; Gold and Harvey, 1993; Blanchard and Neale, 1994; Berman *et al.*, 1997; Basso *et al.*, 1998; Cavadini *et al.*, 2002a). For example, the Wisconsin Card Sorting Test (Heaton, 1981; Feldstein *et al.*, 1999), a somewhat selective test of dorsolateral precortical functioning, typically has been shown to be impaired in schizophrenic patients (Berman *et al.*, 1986; 1992; 1995; Weinberger *et al.*, 1986; 1988; Braff *et al.*, 1991; Marenco *et al.*, 1993; Koren *et al.*, 1998; Saoud *et al.*, 1998) and in patients with frontal lobe damage (Anderson *et al.*, 1991). The Gambling Task (Bechara *et al.*, 1994), with a regional specificity for the ventromedial prefrontal cortex (Bechara *et al.*, 1998; 2000), has produced deficits in psychiatric populations (Abbruzzese *et al.*, 1995), including obsessive-compulsive patients (Cavadini *et al.*, 2002b), patients with antisocial personality disorder (Mazas *et al.*, 2000), individuals with substance abuse disorders (Bartzokis *et al.*, 2000; Grant *et al.*, 2000; Bechara *et al.*, 2001), as well as schizophrenic patients (Wilder *et al.*, 1998). Recently, in an investigation of schizophrenic patients and comparable controls, Ritter *et al.* (2004) measured regional differences in frontal

cortical, *i.e.*, ventromedial prefrontal cortex versus dorsolateral prefrontal cortex, functioning by applying the Wisconsin Card Sorting Test in comparison with the Gambling Task. They found that the schizophrenic patients ( $n=20$ ) performed worse than controls ( $n=15$ ) on the Gambling Task but that no differences were obtained on the Wisconsin Card Sorting Test. Once again, the central influence of decision-making in schizophrenia is to be noted (*cf.*, Carpenter *et al.*, 2000). Since the Gambling Task is considered to reflect ventromedial prefrontal cortex function it appears that in this particular instance that the patients' deficits were restricted to this region only. Nevertheless, it is possible that limitations of this study failed to replicate the volume of previous data. Whatever the case, the application of dual-performance cognitive-task criteria approaches (see also, Beninger, 2006) offers great scope for deriving etiological aspects of comorbidity, both within-task, *i.e.*, instances where two or more disorders share similar deficits, and within-disorder, *i.e.*, instances where two or more tasks that reflect different functional specializations, *e.g.*, executive function versus decision-making, contribution to impairments within the same disorder/brain region. These phenomena, which may be examined at top-down levels of brain activity, constitute further neuronal substrates in comorbidity.

Cognitive impairments are certainly not limited to schizophrenia, a major instance of comorbid overlap is offered by markedly impaired cognitive abilities (Loberg and Miller, 1986; Grant I, 1987; Bolla *et al.*, 2000; Duka *et al.*, 2003), as expressed by problems in visuospatial abstracting, problem-solving, planning and organisation, learning new concepts/notions, flexibility in cognition, selective attention, abstract thinking and various specialized (more-or-less) memory skills, many of these reflecting executive functions (Ardila *et al.*, 1991; Glen *et al.*, 1988; Hambridge, 1990; Waugh *et al.*, 1989; but see also Palomo *et al.*, 2004). These deficits in cognitive functioning are associated with: limited treatment participation, elevated 'drop-out' rates, less after-care utilization and more post-treatment unemployment (Gregson and Taylor, 1977; Donovan *et al.*, 1984; Mackenzie *et al.*, 1987; Teichner *et al.*, 2002), and these individuals are less likely to obtain treatment benefits compared to unimpaired individuals, new to failure to acquire

and integrate new information (Leber *et al.*, 1985; Teichner *et al.*, 2001; Aharonvich *et al.*, 2003). Application of mini-mental state examination both recruits abusers yet may be afflicted by participant exclusion due to cognitive deficits (Crum *et al.*, 1993; Cummings, 1993), thereby endorsing notions of substance abuse-dysfunctional cognition comorbidity (Miller and Saucedo, 1983; O'Farrell and Langenbucher, 1985).

In addressing incentive-learning notions, Schmidt and Beninger (2006) imply that enhanced dopaminergic transmission attributes incentive salience to contextual stimuli that reinforce behaviour, processes underlying psychostimulant-induced context-dependent sensitization contributing to addiction, dyskinesia, amphetamine-induced psychosis and psychostimulant-induced aggression. In certain respects the presentation of an unusual, perhaps counter-intuitive, demonstration of drug-induced behavioural sensitisation, namely haloperidol-induced dopamine-activity reduction leading to behavioural inhibition (not activation as in conventional sensitisation and yet fulfilling all the criterion involving classification of the sensitisation process), provided an astonishing manifestation of neuroplasticity. Remarkably, either a low dose of haloperidol or partial dopamine-depletion (50%) with 6-hydroxydopamine led to the gradual development of catalepsy with repeated testing. Further testing without drug in the case of haloperidol induced an extinction-like stepwise reduction. This highly persistent sensitisation response was shown to be susceptible only to contextual conditions; it was markedly context-dependent. These results would appear to open new horizons implicating neuroplasticity normally associated with neuropsychiatric conditions and symptomatology in neurodegenerative disorders such as Parkinson's disease. From a different, yet comorbid, perspective, the demonstration of a neostriatal habit learning system in amnesic patients and nondemented patients with Parkinson's disease implicates the role of the neostriatum (Knowlton *et al.*, 1996), and provides a parallel to the phenomena treated above. This region, the neostriatum, influences motor behaviour and motor learning, as well as acquisition of non-motor dispositions and tendencies, reliant on novel (contextual) associations, essential for incremental associative learning situations, whereby an intrinsic

neuronal substrate must consideration.

## IMPLICATIONS OF COMORBIDITY

From a perspective of neurologic-psychiatric comorbidity, the Braakian notions of staging in the development of neuropathology (*cf.*, Braak *et al.*, 1994; 2000; 2003; Braak and Del Tredici, 2004), directly obvious in the neurodegenerative disorders but suitable for considerations of neuropsychiatric conditions, have contributed enormously to a necessary paradigmatic shift in the interpretation of wide-ranging observations. Suffice it to say, these standpoints, supported by multiple aspects of gene-environmental interactions, have opened a plethora of both perceivable and as yet vaguely discernible phenomena to further endow current understandings of brain disease states.

## References

- Abbruzzese M, L Bellodi, S Ferri and S Scarone (1995) Frontal lobe dysfunction in schizophrenia and obsessive-compulsive disorder: a neuropsychological study. *Brain Cogn.* **27**, 202-212.
- Aber KM, MP Allin, K Kucharska-Pietura, C Andrew, S Williams, AS David and ML Phillips (2003) Ketamine and fMRI BOLD signal: distinguishing between effects mediated by change in blood flow versus change in cognitive state. *Hum. Brain Map* **18**, 135-145.
- Abercrombie ED and BL Jacobs (1987) Single-unit response of noradrenergic neurons in the locus coeruleus of freely moving cats. I. Acutely presented stressful and nonstressful stimuli. *J. Neurosci.* **7**, 2837-2843.
- Aharonovich E, E Nunes and D Hasin (2003) Cognitive impairment, retention and abstinence among cocaine abusers in cognitive-behavioural treatment. *Drug Alcohol Depend.* **71**, 207-211.
- Akbarian S, WE Bunney Jr, SG Potkin, SB Wigal, JO Hagman, CA Sandman and EG Jones (1993) Altered distribution of nicotinamide-adenine dinucleotide phosphate-diaphorase cells in frontal lobe of schizophrenics implies disturbances of cortical development. *Arch. Gen. Psychiatry* **50**, 169-177.
- Aloise F, G Penna, J Cerase, B Menendez Iglesias and L Adorini (1997) IL 12 production by central nervous system microglia is inhibited by astrocytes. *J. Immunol.* **159**, 1604-1612.
- Alonso M, LR Bevilaqua, I Izquierdo, JH Medina and M Cammarota (2003) Memory formation requires p38MAPK activity in the rat hippocampus. *Neuroreport* **14**, 1989-1992.
- Anderson SW, H Damasio, RD Jones and D Tranel (1991) Wisconsin CARD Sorting Test performance as a measure of frontal lobe damage. *J. Clin. Exp. Neuropsychol.* **13**, 909-922.
- Ardila A, M Roselli and S Strumwasser (1991) Neuropsychological deficits in chronic cocaine abusers. *Int. J. Neurosci.* **57**, 73-79.
- Austin MP, P Mitchell and GM Goodwin (2001) Cognitive deficits in depression: possible implications for functional neuropathology. *Br. J. Psychiatry* **178**, 200-206.
- Aylward RL and S Totterdell (1993) Neurons in the ventral subiculum, amygdala and entorhinal cortex which project to the nucleus accumbens: their input from somatostatin-immunoreactive boutons. *J. Chem. Neuroanat.* **6**, 31-42.
- Bartzokis G, PH Lu, M Beckson, R Rapaport, S Grant, EJ Wiseman and ED London (2000) Abstinence from cocaine reduces high-risk gambling responses on a Gambling Task. *Neuropsychopharmacol.* **22**, 102-103.
- Basso MR, HA Nasrallah, SC Olson and RA Bornstein (1998) Neuropsychological correlates of negative, disorganised and psychotic symptoms in schizophrenia. *Schizophr. Res.* **31**, 99-111.
- Bechara A, AR Damasio, H Damasio and SW Anderson (1994) Insensitivity to future consequences following damage to prefrontal cortex. *Cognition* **50**, 7-15.
- Bechara A, H Damasio, D Tranel and SW Anderson (1998) Dissociation of working memory from decision-making within the human prefrontal cortex. *J. Neurosci.* **18**, 428-437.
- Bechara A, D Tranel and H Damasio (2000) Characterization of the decision-making deficit of patients with ventromedial prefrontal cortex lesions. *Brain* **123**, 2189-2202.
- Bechara A, S Dolan, N Denburg, A Hinds, SW Anderson and PE Nathan (2001) Decision-making deficits, linked to a dysfunctional ventromedial prefrontal cortex, revealed in alcohol and stimulant abusers. *Neuropsychologia* **39**, 376-389.
- Beninger RJ (2006) Dopamine and incentive learning: a framework for considering antipsychotic medication effects. *Neurotox. Res.* **10**, 199-210.
- Berman DE and Y Dudai (2001) Memory extinction, learning anew, and learning the new: dissociations in the molecular machinery of learning in cortex. *Science* **291**, 2417-2419.
- Berman I, B Viegner, A Mason, E Allan, D Pappas and AI Green (1997) Differential relationships between positive and negative symptoms and neuropsychological deficits in schizophrenia. *Schizophr. Res.* **25**, 1-10.
- Berman KF, RF Zec and DR Weinberger (1986) Physiological dysfunction of dorsolateral prefrontal cortex in schizophrenia: II. Role of neuroleptic treatment, attention, and mental effort. *Arch. Gen. Psychiatry* **43**, 126-135.
- Berman KF, EF Torrey, DG Daniel and DR Weinberger (1992) Regional cortical blood flow in monozygotic twins discordant and concordant for schizophrenia. *Arch. Gen. Psychiatry* **49**, 927-934.
- Berman KF, JL Osrem, C Randolph, J Gold, TE Goldberg, R Coppola, RE Carson, P Herscovitch and DR Weinberger (1995) Physiological activation of cortical networks during performance of the Wisconsin Card Sorting Test: a positron emission tomography study. *Neuropsychologia* **33**, 1027-1046.
- Bernabeu R, M Levi de Stein, C Fin, I Izquierdo and JH Medina (1995) Role of hippocampal NO in the acquisi-

- tion and consolidation of inhibitory avoidance learning. *Neuroreport* **6**, 1498-1500.
- Bevilaqua LR, JS Bonini, JI Rossato, LA Izquierdo, M Cammarota and I Izquierdo (2006) The entorhinal cortex plays a role in extinction. *Neurobiol. Learn. Mem.* **85**(2), 192-1977. Epub 2005 Nov 14.
- Bianchin M, RC Da Silva, PK Schmitz, JH Medina and I Izquierdo (1994) Memory of inhibitory avoidance in the rat is regulated by glutamate metabotropic receptors in the hippocampus. *Behav. Pharmacol.* **5**, 356-359.
- Blanchard JJ and JM Neale (1994) The neuropsychological signature of schizophrenia: generalized or differential deficit? *Am. J. Psychiatry* **151**, 40-48.
- Bolla KI, FR Funderburk and JL Cadet (2000) Differential effects of cocaine and concaine alcohol on neurocognitive performance. *Neurology* **45**, 2285-2292.
- Bolton D, P Raven, R Madronal-Luque and IM Marks (2000) Neurological and neuropsychological signs in obsessive compulsive disorder: interaction with behavioural treatment. *Behav. Res. Ther.* **38**, 695-708.
- Braak H and K Del Tredici (2004) Alzheimer's disease: intraneuronal alterations precede insoluble amyloid- $\beta$  formation. *Neurobiol. Aging* **25**, 713-718.
- Braak H, E Braak and EM Mandelkow (1994) A sequence of cytoskeleton changes related to the formation of neurofibrillary tangles and neurophil threads. *Acta Neuropathol.* **87**, 554-567.
- Braak H, K Del Tredici and E Braak (2000) Spectrum of pathology, In: *Mild Cognitive Impairment* (Peterson R, Ed.) (Oxford University Press:New York, NY), pp 149-189.
- Braak H, KD Del Tredici, U Rüb, RAI de Vos, ENH Jansen Steur and E Braak (2003) Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol. Aging* **24**, 197-211.
- Braff DL, R Heaton, J Kuck, M Cullum, J Moranville, I Grant and S Zisook (1991) The generalized pattern of neuropsychological deficits in outpatients with chronic schizophrenia with heterogeneous Wisconsin Card Sorting Test results. *Arch. Gen. Psychiatry* **48**, 891-898.
- Brown AS, MD Begg, S Gravenstein, CA Schaefer, RJ Wyatt, M Bresnahan, VP Babulas and ES Susser (2004) Serologic evidence of prenatal influenza in the etiology of schizophrenia. *Arch. Gen. Psychiatry* **61**, 774-780.
- Buchanan RW, K Vldar, PE Barta and GD Pearlson (1998) Structural evaluation of the prefrontal cortex in schizophrenia. *Am. J. Psychiatry* **155**, 1049-1055.
- Buka SL, MT Tsuang, EF Torrey, MA Klebanoff, D Berstein and RH Yolken (2001) Maternal infections and subsequent psychosis in the offspring. *Arch. Gen. Psychiatry* **58**, 1032-1037.
- Callaway CW, RL Hakan and SJ Henriksen (1991) Distribution of the amygdala input to the nucleus accumbens septi: an electrophysiological investigation. *J. Neural Transm. Gen. Sect.* **83**, 215-225.
- Cammarota M, LR Bevilaqua, JI Rossato, C Zinn, JH Medina and I Izquierdo (2007) MAPK and CaMKII-mediated events during memory formation, In: *Comorbidity in Neuropsychiatric Disorders* (Palomo T, RJ Beninger, RM Kostrzewa and T Archer, Eds.) (Fundacion Cerebro y Mente: Madrid) (In press).
- Campodónico JR, EH Aylward, AM Codori, C Young, L Krafft, M Magdalinski, N Ranen, P Slavney and J Brandt (1998) When does Huntington's disease begin? *J. Int. Neuropsychol. Soc.* **4**, 467-473.
- Carlsson A, N Waters, S Holm-Waters, J Tedroff, M Nilsson and ML Carlsson (2001) Interactions between monoamines, glutamate and GABA in schizophrenia: new evidence. *Ann. Rev. Pharmacol. Toxicol.* **41**, 237-260.
- Carpenter WT, JM Gold, AC Lahti, CA Queern, RR Conley, JJ Bartko, J Kovnick and PS Appelbaum (2000) Decisional capacity for informed consent in schizophrenia research. *Arch. Gen. Psychiatry* **57**, 533-538.
- Carr DB, P O'Donnell, JP Card and SR Sesack (1999) Dopamine terminals in the rat prefrontal cortex synapse on pyramidal cells that project to the nucleus accumbens. *J. Neurosci.* **19**, 11049-11060.
- Caspi K, K Sugden, TE Moffitt, A Taylor, IW Craig, H Harrington, J McClay, J Mill, A Martin, A Braithwaite and R Poulton (2003) Influence of life stress on depression: moderation by a polymorphism in the 5.THH gene. *Science* **301**, 386-389.
- Cavedini P, R Giovanna, R Keller, A D'Annuncci and L Bellodi (2002a) Frontal lobe dysfunction in pathological gambling patients. *Biol. Psychiatry* **51**, 334-341.
- Cavedini P, G Riboldi, A D'Annuncci, P Belotti, M Cisima and L Bellodi (2002b) Decision-making heterogeneity in obsessive-compulsive disorder: ventromedial prefrontal cortex function predicts different treatment outcomes. *Neuropsychologia* **40**, 205-211.
- Chambers RA, JH Krystal and DW Self (2001) A neurobiological basis for substance abuse comorbidity in schizophrenia. *Biol. Psychiatry* **50**, 71-83.
- Chambers RA, JR Taylor and MN Potenza (2003) Developmental neurocircuitry of motivation in adolescence: a critical period of addiction vulnerability. *Am. J. Psychiatry* **160**, 1041-1052.
- Coco ML, CM Kuhn, TD Ely and CD Kilts (1992) Selective activation of mesoamygdaloid dopamine neurons by conditioned stress: attenuation by diazepam. *Brain Res.* **590**, 39-47.
- Colder CR and L Chassin (1993) The stress and negative affect model of adolescent alcohol use and the moderating effects of behavioural undercontrol. *J. Stud. Alcohol* **54**, 326-333.
- Collier DA and T Li (2003) The genetics of schizophrenia: glutamate not dopamine. *Eur. J. Pharmacol.* **480**, 177-184.
- Cook GS and AV Hill (2001) Genetics of susceptibility to human infectious diseases. *Nat. Rev. Genetics* **2**, 967-977.
- Correll CM, JA Rosenkranz and AA Grace (2005) Chronic cold stress alters prefrontal cortical modulation of amygdala neuronal activity in rats. *Biol. Psychiatry* **58**, 382-391.
- Coryell W, P Nopoulos, W Drevets, T Wilson and NC Andreasen (2005) Subgenual prefrontal cortex volumes in major depressive disorder and schizophrenia: diagnostic specificity and prognostic implications. *Am. J. Psychiatry* **162**, 1706-1712.
- Crombag HS, G Gorny, Y Li, B Kolb and TE Robinson (2005) Opposite effects of amphetamine self-administration experience on dendritic spines in the medial and orbital prefrontal

- cortex. *Cereb. Cortex* **15**, 341-348.
- Crum RM, JC Anthony, SS Bassett and MF Folstein (1993) Population-based norms for the mini-mental state examination by age and educational level. *JAMA* **269**, 2386-2391.
- Cummings JL (1993) Mini-mental state examination: norms, normals and numbers. *JAMA* **269**, 2420-2421.
- Davidson RJ, DC Jackson and NH Kalin (2000) Emotion, plasticity, context, and regulation: perspectives from affective neuroscience. *Psychol. Bull.* **126**, 890-909.
- Davidson RJ, DA Lewis, LB Alloy, DG Amaral, G Bush, JD Cohen, WC Drevets, MJ Farah, J Kagan, JL McClelland, S Nolen-Hoeksema and BS Peterson (2002) Neural and behavioral substrates of mood and mood regulation. *Biol. Psychiatry* **52**, 478-502.
- Dierker LC, S Avenevoli, M Stolar and KR Merikangas (2002) Smoking and depression: an examination of mechanisms of comorbidity. *Am. J. Psychiatry* **159**, 947-953.
- Ding DC, PL Gabbott and S Totterdell (2001) Differences in the laminar origin of projections from the medial prefrontal cortex to the nucleus accumbens shell and core regions in the rat. *Brain Res.* **917**, 81-89.
- Donovan DM, DR Kivlahan and RD Walker (1984) Clinical limitations of neuropsychological testing in predicting treatment outcome among alcoholics. *Alcohol Clin. Exp. Res.* **8**, 470-475.
- Drevets WC (2000) Neuroimaging studies of mood disorders. *Biol. Psychiatry* **48**, 813-829.
- Dudai Y (2002) Molecular bases of long-term memories: a question of persistence. *Curr. Opin. Neurobiol.* **12**, 211-216.
- Dudai Y (2004) The neurobiology of consolidations, or, how stable is the engram? *Annu. Rev. Psychol.* **55**, 51-86.
- Duka T, JM Townshend, K Collier and DN Stephens (2003) Impairment in cognitive function after multiple detoxifications in alcoholic inpatients. *Alcohol Clin. Exp. Res.* **27**, 1563-1572.
- Dunn AJ (1988) Stress-related activation of cerebral dopaminergic systems. *Ann. NY Acad. Sci.* **537**, 188-205.
- Eastwood SL and PJ Harrison (2001) Synaptic pathology in the anterior cingulate cortex in schizophrenia and mood disorders. A review and a Western blot study of synaptophysin, GAP-43 and the complexins. *Brain Res. Bull.* **55**, 569-578.
- Eastwood SL, PW Burnet, R Gittins, K Baker and PJ Harrison (2001) Expression of serotonin 5-HT<sub>2A</sub> receptors in the human cerebellum and alterations in schizophrenia. *Synapse* **42**, 104-114.
- Ebert D and KP Ebmeier (1996) The role of the cingulate gyrus in depression: from functional anatomy to neurochemistry. *Biol. Psychiatry* **39**, 1044-1050.
- Ebmeier KP and D Kronhaus (2002) Brain imaging and mood disorders. In: *Biological Psychiatry* (D'haenen H, JA den Boer and P Willner, Eds.) (Wiley & sons: London).
- Ebmeier KP, E Rose E and D Steele (2006) Cognitive impairment and fMRI in major depression. *Neurotox. Res.* **10**(2), 87-92.
- Elliott R, KJ Friston and RJ Dolan (2000) Dissociable neural responses in human reward systems. *J. Neurosci.* **20**, 6159-6165.
- Elliott R, JS Rubensztein, BJ Sahakian and RJ Dolan (2002) The neural basis of mood-congruent processing biases in depression. *Arch. Gen. Psychiatry* **59**, 597-604.
- Espey MG, ON Chemshev, JJ Reinhard, MA Namboodiri and CA Colton (1997) Activated human microglia produce the excitotoxin quinolinic acid. *Neuroreport* **8**, 431-434.
- Feigin AS, Y Ma, MF Ghilardi, M Fukuda, SM Ivanova, CP Ghez and D Eidelberg (2001) Abnormal thalamocortical activation responses during motor sequence learning in pre-symptomatic Huntington's disease. *Soc. Neurosci. Abstr.* **27**.
- Feldstein SN, FR Keller, RE Portman, RL Durham, KJ Klebe and PD Hasker (1999) A comparison of computerised and standard versions of the Wisconsin Card Sorting Test. *Clin. Neuropsychol.* **13**, 303-313.
- Finlay JM, MJ Zigmond and ED Abercrombie (1995) Increased dopamine and norepinephrine release in medial prefrontal cortex induced by acute and chronic stress: effects of diazepam. *Neuroscience* **64**, 619-628.
- Finlay JM, HP Jedema, AD Rabinovic, MJ Mana, MJ Zigmond and AF Sved (1997) Impact of corticotropin-releasing hormone on extracellular norepinephrine in prefrontal cortex after chronic cold stress. *J. Neurochem.* **69**, 144-150.
- Fornai F, MG Alessandri, MT Torracca, L Bassi and GU Corsini (1997) Effects of noradrenergic lesions on MPTP/MPP<sup>+</sup> kinetics and MPTP-induced nigrostriatal dopamine depletions. *J. Pharmacol. Exp. Ther.* **283**, 100-107.
- Fossati P, F Coyette, AM Ergis and JF Allilaire (2002) Influence of age and executive functioning on verbal memory of inpatients with depression. *J. Affect. Disord.* **68**, 261-271.
- Fossati P, PO Harvey, G Le Bastard, AM Ergis, R Jouvent and JF Allilaire (2004) Verbal memory performance of patients with a first depressive episode and patients with unipolar and bipolar recurrent depression. *J. Psychiatr. Res.* **38**, 137-144.
- Fredriksson A and T Archer (2006) Subchronic administration of haloperidol influences the functional deficits of postnatal iron administration in mice. *Neurotox. Res.* **10**(2), 123-129.
- French SJ and S Totterdell (2002) Hippocampal and prefrontal cortical inputs monosynaptically converge with individual projection neurons of the nucleus accumbens. *J. Comp. Neurol.* **446**(2), 151-165.
- French SJ and S Totterdell (2003) Individual nucleus accumbens-projection neurons receive both basolateral amygdala and ventral subicular afferents in rats. *Neuroscience* **119**, 19-31.
- Gehring WJ and AR Willoughby (2002) The medial frontal cortex and the rapid processing of monetary gains and losses. *Science* **295**, 2279-2282.
- George TP and JH Krystal (2000) Comorbidity of psychiatric and substance abuse disorders. *Curr. Opin. Psychiatry* **13**, 327-331.
- Gilder DA, TL Wall and CL Ehlers (2004) Comorbidity of select anxiety and affective disorders with alcohol dependence in Southwest California Indians. *Alcohol Clin. Exp. Res.* **28**, 1805-1813.
- Glantz LA and DA Lewis (2000) Decreased dendritic spine density on prefrontal cortical neurons in schizophrenia. *Arch. Gen. Psychiatry* **57**, 65-73.
- Glenn SW, OA Parsons, R Sinha and L Stevens (1988) The effects of repeated withdrawals from alcohol on the memory of male and female alcoholics. *Alcohol Alcohol.* **23**, 337-

- 342.
- Gold JM and PD Harvey (1993) Cognitive deficits in schizophrenia. *Psychiatr. Clin. North Am.* **16**, 295-312.
- Goldberg TE, JR Kelson, DR Weinberger, NH Pliskin, PD Kirwin and KF Berman (1988) Performance of schizophrenic patients on putative neuropsychological tests of frontal lobe function. *Int. J. Neurosci.* **42**, 51-58.
- Goldberg TE, JD Ragland, EF Torrey, JM Gold, LB Bigelow and DR Weinberger (1990) Neuropsychological assessment of monozygotic twins discordant for schizophrenia. *Arch. Gen. Psychiatry* **47**, 1066-1072.
- Goldberg TE, EF Torrey, JM Gold, JD Ragland, LB Bigelow and DR Weinberger (1993) Learning and memory in monozygotic twins discordant for schizophrenia. *Psychol. Med.* **23**, 71-85.
- Golik-Gruber V, D Brietenfeld, EN Gruber and D Karlovic (2003) Sulpiride psychopharmacotherapy in patients with alcohol addiction and depression comorbidity. *Acta Clinica Croatia* **42**, 29-33.
- Goto Y and AA Grace (2005) Dopamine-dependent interactions between limbic and prefrontal cortical synaptic plasticity in the nucleus accumbens: disruption by cocaine sensitization. *Neuron* **47**, 255-266.
- Goto Y and Grace AA (2006) Alterations in medial prefrontal cortical activity and plasticity in rats with disruption of cortical development. *Biol. Psychiatry* 2006 Aug 31; [Epub ahead of print]
- Grace AA (2006) Disruption of cortical-limbic interaction as a substrate for comorbidity. *Neurotox. Res.* **10**(2), 93-101.
- Grace AA, H Moore and P O'Donnell (1998) The modulation of corticoaccumbens transmission by limbic afferents and dopamine: a model for the pathophysiology of schizophrenia. *Adv. Pharmacol.* **42**, 721-724.
- Grant BF (1995) Comorbidity between DSM-IV drug use disorders and major depression: results of a national survey. *J. Subst. Abuse* **7**, 481-497.
- Grant I (1987) Alcohol and the brain: neuropsychological correlates. *J. Consult. Clin. Psychol.* **55**, 310-324.
- Grant S, C Contoreggi and ED London (2000) Drug abusers show impaired performance in a laboratory test of decision-making. *Neuropsychologia* **38**, 1180-1187.
- Gray TS (1999) Functional and anatomical relationships among the amygdala, basal forebrain, ventral striatum, and cortex. An integrative discussion. *Ann. NY Acad. Sci.* **877**, 439-444.
- Gregson RAM and GM Taylor (1977) Prediction of relapse in men alcoholics. *J. Stud. Alcohol* **38**, 1749-1760.
- Gresch PJ, AF Sved, MJ Zigmond and JM Finlay (1994) Stress-induced sensitization of dopamine and norepinephrine efflux in medial prefrontal cortex of the rat. *J. Neurochem.* **63**, 575-583.
- Groenewegen HJ, Y Galis-de Graaf and WJ Smeets (1999) Integration and segregation of limbic cortico-striatal loops at the thalamic level: an experimental tracing study in rats. *J. Chem. Neuroanat.* **16**(3), 167-185.
- Hambrecht M and H Hafner (2000) Cannabis, vulnerability, and the onset of schizophrenia: an epidemiological perspective. *Aust. NZ J. Psychiatry* **34**, 468-475.
- Hambridge DM (1990) Intellectual impairment in male alcoholics. *Alcohol Alcohol.* **25**, 555-559.
- Hariri AR, VS Mattay, A Tessitore, F Fera and DR Weinberger (2003) Neocortical modulation of the amygdala response to fearful stimuli. *Biol. Psychiatry* **53**, 494-501.
- Harrison PJ, AJ Law and SL Eastwood (2003) Glutamate receptors and transporters in the hippocampus in schizophrenia. *Ann. NY Acad. Sci.* **1003**, 94-101.
- Harvey PO, G Le Bastard, JB Pochon, R. Levy, JF Allilaire, B Dubois and P Fossati (2004) Executive functions and updating of the contents of working memory in unipolar depression. *J. Psychiatr. Res.* **38**, 567-576.
- Heaton RK (1981) *Wisconsin Card Sorting Test Manual* (PAR: Odessa, FL).
- Higihara A, K Tarumi and K Nobutomo (2003) Positive and negative effects of social support on the relationship between work stress and alcohol consumption. *J. Stud. Alcohol* **64**, 874-883.
- Hill AV (1999) Genetics of infectious disease resistance. *Curr. Opin. Genet. Dev.* **6**, 348-353.
- Izquierdo I and JH Medina (1995) Correlation between the pharmacology of long-term potentiation and the pharmacology of memory. *Neurobiol. Learn. Mem.* **63**, 19-32.
- Izquierdo I and JH Medina (1997) Memory formation: the sequence of biochemical events in the hippocampus and its connection to activity in other brain structures. *Neurobiol. Learn. Mem.* **68**, 285-316.
- Izquierdo I, LRM Bevilaqua, JI Rossato, JS Bonini, WC Da Silva, JH Medina and M Cammarota (2006) The connection between the hippocampal and the striatal memory systems of the brain: a review of recent findings. *Neurotox. Res.* **10**(2), 113-121.
- Jedema HP, AF Sved, MJ Zigmond and JM Finlay (1999) Sensitization of norepinephrine release in medial prefrontal cortex: effect of different chronic stress protocols. *Brain Res.* **830**, 211-217.
- Jedema HP, JM Finlay, AF Sved and AA Grace (2001) Chronic cold exposure potentiates CRH-evoked increases in electrophysiological activity of locus coeruleus neurons. *Biol. Psychiatry* **49**, 351-359.
- Johnson JG, P Cohen, DS Pine, DF Klein, S Kasen and JS Brook (2000) Association between cigarette smoking and anxiety disorders during adolescence and early childhood. *JAMA* **248**, 2348-2351.
- Johnson LR, RL Aylward, Z Hussain and S Totterdell (1994) Input from the amygdala to the rat nucleus accumbens: its relationship with tyrosine hydroxylase immunoreactivity and identified neurons. *Neuroscience* **61**, 851-865.
- Katarina Z, K Andrew, M Filomena and H Xu-Feng (2004) Investigation of m1/m4 muscarinic receptors in the anterior cingulate cortex in schizophrenia, bipolar disorder, and major depression disorder. *Neuropsychopharmacology* **29**, 619-625.
- Kauer JA (2004) Learning mechanisms in addiction: synaptic plasticity in the tegmental area as a result of exposure to drugs of abuse. *Ann. Rev. Physiol.* **66**, 447-475.
- Kendler KS, AC Heath, MC Neale, RC Kessler and LJ Eavea (1993) Alcoholism and major depression in women. A twin study of the causes of comorbidity. *Arch. Gen. Psychiatry* **50**, 690-698.

- Keuthen NJ, RL O'Sullivan and S Sprich-Buckminster (1998) Trichotillomania: current issues in conceptualization and treatment. *Psychother. Psychosom.* **67**, 202-213.
- Knowlton BJ, JA Mangels and LR Squire (1996) A neostriatal habit learning system in humans. *Science* **273**, 1399-1402.
- Koponen H, P Rantakallio, J Veijola, P Jones, J Jokelainen and M Isohanni (2004) Childhood central nervous system infections and risk for schizophrenia. *Eur. Arch. Psychiatry Clin. Neurosci.* **254**, 9-13.
- Koren D, WS Harrison, WS Kremen, JM Goldstein, SV Faraone, LJ Seidman, MJ Lyons, B Caplan and MT Tsuang (1998) Factor structure of the Wisconsin Card Sorting Test: dimensions of deficit in schizophrenia. *Neuropsychology* **12**, 289-302.
- Kornhuber J, J Wiltfang and S Bleich (2004) The etiopathogenesis of schizophrenia. *Pharmacopsychiatry* **37** (Suppl. 2), S103-S112.
- Kroner S, JA Rosenkranz, AA Grace and G Barrionuevo (2005) Dopamine modulates excitability of basolateral amygdala neurons *in vitro*. *J. Neurophysiol.* **93**, 1598-1610.
- Lavolette SR, WJ Lipski and AA Grace (2005) A subpopulation of neurons in the medial prefrontal cortex encode emotional learning with burst and frequency codes through a dopamine D<sub>4</sub> receptor-dependent basolateral amygdala input. *J. Neurosci.* **25**, 6066-6075.
- Le Doux JE (2000) Emotion circuits in the brain. *Ann. Rev. Neurosci.* **23**, 155-184.
- Leber WR, OR Parsons and N Nichols (1985) Neuropsychological test results are related to ratings of men alcoholics' therapeutic progress: a replicated study. *J. Stud. Alcohol* **46**, 116-121.
- Leonard BE and A Myint (2006) Inflammation and depression: is there a causal connection with dementia? *Neurotox. Res.* **10**(2), 149-160.
- Lewis DA, DW Volk and T Hashimoto (2004) Selective alterations in prefrontal cortical GABA neurotransmission in schizophrenia: a novel target for the treatment of working memory dysfunction. *Psychopharmacology* **174**, 143-150.
- Loberg T and WR Miller (1986) Personality, cognitive and neuropsychological correlates of harmful alcohol consumption: a cross-national comparison of clinical samples. *Ann. NY Acad. Sci.* **472**, 75-97.
- Lochner C, S Seedat, PL du Toit, DG Nel, DJ Niehaus, R Sandler and DJ Stein (2005) Obsessive-compulsive disorder and trichotillomania: a phenomenological comparison. *BMC Psychiatry* **5**, 2.
- Mackenzie A, FR Funderburk, RP Allen and RL Stefan (1987) The characteristics of alcoholics frequently lost to follow-up. *J. Stud. Alcohol* **48**, 119-123.
- Mana MJ and AA Grace (1997) Chronic cold stress alters the basal and evoked electrophysiological activity of rat locus coeruleus neurons. *Neuroscience* **81**, 1055-1064.
- Maren S and GJ Quirk (2004) Neuronal signalling of fear memory. *Nat. Rev. Neurosci.* **5**, 844-852.
- Marenco S, R Coppola, DG Daniel, JR Zigun and DR Weinberger (1993) Regional cerebral blood flow during the Wisconsin Card Sorting Test in normal subjects as studied by xenon-133 dynamic SPECT: comparison of absolute values, percent distribution values, and covariance analysis. *Psychiatry Res.* **50**, 177-192.
- Marien MR, M Briley and FC Colpaert (1993) Noradrenaline depletion exacerbates MPTP-induced striatal dopamine loss in mice. *Eur. J. Pharmacol.* **236**, 487-489.
- Mayberg HS (1997) Limbic-cortical dysregulation: a proposed model of depression. *J. Neuropsychiatry Clin. Neurosci.* **9**, 471-481.
- Mayberg HS (2003) Modulating dysfunctional limbic-cortical circuits in depression: towards development of brain-based algorithms for diagnosis and optimised treatment, In: *Imaging Neuroscience: Clinical Frontiers for Diagnosis and Management* (Frackowiak RS and T Jones, Eds.) (Oxford University Press: Oxford), pp 193-207.
- Mazas C, PR Finn and JE Steinmetz (2000) Decision-making biases, antisocial personality, and early-onset alcoholism. *Alcohol Clin. Exp. Res.* **24**, 1036-1040.
- McClure SM, MK York and PR Montague (2004) The neural substrates of reward processing in humans: the modern role of fMRI. *Neuroscientist* **10**, 260-268.
- McGaugh JL (2000) Memory - a century of consolidation. *Science* **287**, 248-251.
- McGaugh JL and B Roozendaal (2002) Role of adrenal stress hormones in forming lasting memories in the brain. *Curr. Opin. Neurobiol.* **12**, 205-210.
- Medina JH, N Schröder and I Izquierdo (1999) Two different properties of short- and long-term memory. *Behav. Brain Res.* **103**, 119-121.
- Mednick SA, RA Machon, MO Huttunen and D Bonett (1988) Adult schizophrenia following prenatal exposure to an influenza epidemic. *Arch. Gen. Psychiatry* **45**, 817-824.
- Milad MR and GJ Quirk (2002) Neurons in medial prefrontal cortex signal memory for fear extinction. *Nature* **420**, 70-74.
- Miller CL, IC Llenos, JR Dulay, MM Barillo, RH Yolken and S Weis (2004) Expression of the kynurenine pathway enzyme tryptophan 2,3-dioxygenase is increased in the frontal cortex of individuals with schizophrenia. *Neurobiol. Dis.* **15**, 618-629.
- Miller WR and CF Saucedo (1983) Assessment of neuropsychological impairment and brain damage in problem drinkers, In: *Clinical Neuropsychology: Interface with Neurologic and Psychiatric Disorders* (Golden CJ, JA Moses, JA Coffman, WR Miller and FD Strider, Eds.) (Grune and Stratton: New York), pp 141-195.
- Modell JG, JM Mountz, GC Curtis and JF Greden (1989) Neurophysiologic dysfunction in basal ganglia/limbic striatal thalamocortical circuits as a pathogenetic mechanism of obsessive-compulsive disorder. *J. Neuropsychiatry Clin. Neurosci.* **1**, 27-36.
- Moore H, HJ Rose and AA Grace (2001) Chronic cold stress reduces the spontaneous activity of ventral tegmental dopamine neurons. *Neuropsychopharmacology* **24**, 410-419.
- Müller N (2004) Immunological and infectious aspects of schizophrenia (Editorial). *Eur. Arch. Psychiatry Clin. Neurosci.* **254**, 1-3.
- Müller N and Schwarz M (2006) The immune function in schizophrenia: an inflammation-mediated dysbalance of the glutamatergic neurotransmission. *Neurotox. Res.* **10**(2),

- 131-148.
- Müller N, M Ackenheil, E Hofschuster, W Mempel and R Eckstrin (1991) Cellular immunity in schizophrenic patients before and during neuroleptic therapy. *Psychiatry Res.* **37**, 147-160.
- Müller N, M Empel, M Riedel, MJ Schwarz and M Ackenheil (1997a) Neuroleptic treatment increases soluble IL-2 receptors and decreases soluble IL-6 receptors in schizophrenia. *Eur. Arch. Psychiatry Clin. Neurosci.* **247**, 308-313.
- Müller N, P Dobmeier, M Empel, M Riedel, MJ Schwarz and M Ackenheil (1997b) Soluble IL-6 receptors in the serum and cerebrospinal fluid of paranoid schizophrenic patients. *Eur. Psychiatry* **12**, 294-299.
- Müller N, M Hadjamu, M Riedel, J Primbs, M Ackenheil and R Gruber (1999) The adhesion-molecule receptor expression on T helper cells increases during treatment with neuroleptics and is related to the blood-brain barrier permeability in schizophrenia. *Am. J. Psychiatry* **156**, 634-636.
- Müller N, M Riedel, M Ackenheil and MJ Schwarz (2000) Cellular and humoral immune system in schizophrenia: a conceptual re-evaluation. *World J. Biol. Psychiatry* **1**, 173-179.
- Müller N, M Riedel, C Scheppach, B Brandstätter, S Sokkullu, K Krampe, M Olmschneider, H-J Möller and M Schwarz (2002) Beneficial anti-psychoic effects of celecoxib add-on therapy compared to risperidone alone in schizophrenia. *Am. J. Psychiatry* **159**, 1029-1034.
- Murphy FC, A Michael, TW Robbins and BJ Sahakian (2003) Neuropsychological impairment in patients with major depressive disorder: the effects of feedback on task performance. *Psychol. Med.* **33**, 455-467.
- Nisenbaum LK, MJ Zigmond, AF Sved and ED Abercrombie (1991) Prior exposure to chronic stress results in enhanced synthesis and release of hippocampal norepinephrine in response to a novel stressor. *J. Neurosci.* **11**, 1478-1484.
- Nishi K, T Kondo and H Narabayashi (1991) Destruction of norepinephrine terminals in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-treated mice reduces locomotor activity induced by L-dopa. *Neurosci. Lett.* **123**, 244-247.
- Numakawa T, Y Yagasaki, T Ishimoto, T Okada, T Suzuki, N Iwata, N Ozaki, T Taguchi, M Tatsumi, K Kamijima, RE Straub, DR Weinberger, H Kunugi and R Hashimoto (2004) Evidence of novel neuronal functions of dysbindin, a susceptibility gene for schizophrenia. *Hum. Mol. Genet.* **13**, 2699-2708.
- O'Donnell P and AA Grace (1995) Synaptic interactions among excitatory afferents to nucleus accumbens neurons: hippocampal gating of prefrontal cortical input. *J. Neurosci.* **15**, 3622-3639.
- O'Farrell TT and JW Langenbucher (1985) Alcohol abuse, In: *Practice of Inpatient Behavior Therapy: a Clinical Guide* (Hersen M, Ed.) (Grune and Stratton:Orlando), pp 141-172.
- O'Sullivan RL, CS Mansueto, EA Lerner and EC Miguel (2000) Characterization of trichotillomania. A phenomenological model with clinical relevance to obsessive-compulsive spectrum disorders. *Psychiatr. Clin. North. Am.* **23**, 587-604.
- Palomo T, T Archer, RJ Beninger and RM Kostrzewa (2004) Gene-environment interplay in neurogenesis and neurodegeneration. *Neurotox. Res.* **6**(6), 415-434.
- Pierri JN, AS Chaudry, TU Woo and DA Lewis (1999) Alterations in chandelier neuron axon terminals in the prefrontal cortex of schizophrenic subjects. *Am. J. Psychiatry* **156**, 1709-1719.
- Pochon JB, R Levy, P Fossati, S Lehericy, JB Poline, B Pillon, D Le Bihan and B Dubois (2002) The neural system that bridges reward and cognition in humans: an fMRI study. *Proc. Natl. Acad. Sci. USA* **99**, 5669-5674.
- Porter RJ, P Gallagher, JM Thompson and AH Young (2003) Neurocognitive impairment in drug-free patients with major depressive disorder. *Br. J. Psychiatry* **182**, 214-220.
- Quirk GJ, GK Russo, JL Barron and K Lebron (2000) The role of ventromedial prefrontal cortex in the recovery of extinguished fear. *J. Neurosci.* **20**, 6225-6231.
- Rajkowska G, JJ Miguel-Hidalgo, J Wei, G Dilley, SD Pittman, HY Meltzer, JC Overholser, BL Roth and CA Stockmeier (1999) Morphometric evidence for neuronal and glial prefrontal cell pathology in major depression. *Biol. Psychiatry* **45**, 1085-1098.
- Ritter LM, JH Meador-Woodruff and GW Dalack (2004) Neurocognitive measures of prefrontal cortical dysfunction in schizophrenia. *Schizophr. Res.* **68**, 65-73.
- Rodriguez-Moreno A, M Carrion and JM Delgado-Garcia (2006) The nicotinic agonist RJR-2403 compensates the impairment of eyeblink conditioning produced by the non-competitive NMDA-receptor antagonist MK-801. *Neurosci. Lett.* **402**(1-2), 102-107.
- Rohsenow DJ (1982) Social anxiety, daily moods, and alcohol use over time among heavy social drinking men. *Addict. Behav.* **7**, 311-315.
- Romelsjo A and D Hasin (1992) The relationship between stressful working conditions and high alcohol consumption. *Br. J. Addict.* **87**, 1173-1183.
- Rose EJ, E Simonotto and KP Ebmeier (2005) Limbic overactivity in depression is associated with impaired working memory during the n-back task. *Neuroimage* **29**(1), 203-215. Epub 2005 Sep. 12.
- Rosenkranz JA and AA Grace (2001) Dopamine attenuates prefrontal cortical suppression of sensory inputs to the basolateral amygdala of rats. *J. Neurosci.* **21**, 4090-4103.
- Rosenkranz JA and AA Grace (2002a) Cellular mechanisms of infralimbic and prelimbic prefrontal cortical inhibition and dopaminergic modulation of basolateral amygdala neurons *in vivo*. *J. Neurosci.* **22**, 324-337.
- Rosenkranz JA and AA Grace (2002b) Dopamine-mediated modulation of odour-evoked amygdala potentials during Pavlovian conditioning. *Nature* **417**, 282-287.
- Rosenkranz JA and AA Grace (2003) Affective conditioning in the basolateral amygdala of anesthetized rats is modulated by dopamine and prefrontal cortical inputs. *Ann. NY Acad. Sci.* **985**, 488-491.
- Saoud M, T Hueber, H Mandran, J Dalery and T D'Amato (1998) Olfactory identification deficiency and WCST performance in men with schizophrenia. *Psychiatry Res.* **81**, 251-257.
- Sapolsky RM (1996) Stress, glucocorticoids, and damage to the nervous system: the current state of confusion. *Stress* **1**, 1-19.

- Sapolsky RM (2001) Depression, antidepressants and the shrinking hippocampus. *Proc. Natl. Acad. Sci. USA* **98**, 12320-12322.
- Sapolsky RM (2005) The influence of social hierarchy on primate health. *Science* **308**, 648-652.
- Sapolsky RM, H Uno, CS Rebert and CE Finch (1990) Hippocampal damage associated with prolonged glucocorticoid exposure in primates. *J. Neurosci.* **10**, 2897-2902.
- Schmidt WJ and RJ Beninger (2006) Behavioural sensitization in addiction, schizophrenia, Parkinson's disease and dyskinesia. *Neurotox. Res.* **10**(2), 161-166.
- Schwab SG, M Knapp, S Mondabon, J Hallmayer, M Borrmann-Hassenbach, M Albus, B Lerer, M Rietschel, M Trixler, W Maier and DB Wildenauer (2003) Support for association of schizophrenia with genetic variation in the 6p22.3 gene, dysbindin, in sib-pair families with linkage and in an additional sample of triad families. *Am. J. Hum. Genet.* **72**, 185-190.
- Schwarcz R, A Rassoulpour, HQ Wu, D Medoff, CA Tamminga and RC Roberts (2001) Increased cortical kynurenate content in schizophrenia. *Biol. Psychiatry* **50**, 521-530.
- Schwarz M, S Chiang, N Müller and M Ackenheil (2001) T-Helper-1 and T-Helper-2 responses in psychiatric disorders. *Brain Behav. Immunol.* **15**, 340-370.
- Shah PJ, KP Ebmeier, ME Glabus MF Glabus and GM Goodwin (1998) cortical grey matter reductions associated with treatment-resistant chronic unipolar depression. Controlled magnetic resonance imaging study. *Br. J. Psychiatry* **172**, 527-532.
- Siegle GJ, SR Steinhauer, ME Thase, VA Stenger and CS Carter (2002) Can't shake that feeling: event-related fMRI assessment of sustained amygdala activity in response to emotional information in depressed individuals. *Biol. Psychiatry* **51**, 693-707.
- Smelson DA, MF Losonczy, CW Davis, M Kaune, J Williams and D Ziedonis (2002) Risperidone decreases craving and relapses in individuals with schizophrenia and cocaine dependence. *Can. J. Psychiatry* **47**, 671-675.
- Speciale C and R Schwarcz (1993) On the production and disposition of quinolinic acid in rat brain and liver slices. *J. Neurochem.* **60**, 212-218.
- Spedding M, I Neau and L Harsing (2003) Brain plasticity and pathology in psychiatric disease: sites of action for potential therapy. *Curr. Opin. Pharmacol.* **3**, 33-40.
- Steele JD and SM Lawrie (2004) Segregation of cognitive and emotion function in the prefrontal cortex: a stereotactic meta-analysis. *Neuroimage* **21**, 868-875.
- Steele JD, M Meyer and KP Ebmeier (2004) Neural predictive error signal correlates with depressive illness severity in a game paradigm. *Neuroimage* **23**, 269-280.
- Stefansson H, E Sigurdsson, V Steinthorsdottir, S Björnsdottir, T Sigmundsson, S Ghosh, J Brynjolfsson, S Gunnarsdottir, O Ivarsson *et al.* (2002) Neureglin 1 and susceptibility to schizophrenia. *Am. J. Hum. Genet.* **71**, 877-892.
- Stefansson H, J Sarginson, A Kong, P Yates, V Steinthorsdottir, E Gudfinnsson, S Gunnarsdottir, N Walker, H Petursson *et al.* (2003) Association of neureglin 1 with schizophrenia confirmed in a Scottish population. *Am. J. Hum. Genet.* **72**, 83-87.
- Stein DJ, D Simeon, LJ Cohen and E Hollander (1995) Trichotillomania and obsessive-compulsive disorder. *J. Clin. Psychiatry* **56**, 28-34.
- Stine OC, N Pleasant, ML Franz, MH Abbott, SE Folstein and CA Ross (1993) Correlation between onset age of Huntington's disease and length of the trinucleotide repeat in IT-15. *Hum. Mol. Genet.* **2**, 1547-1549.
- Swedo SE and HL Leonard (1992) Trichotillomania. An obsessive compulsive spectrum disorder? *Psychiatr. Clin. North. Am.* **15**, 777-790.
- Swendsen JD and KR Merikangas (2000) The comorbidity of depression and substance use disorders. *Clin. Psychol. Rev.* **20**, 173-189.
- Szechtman H and EZ Woody (2004) Obsessive-compulsive disorder as a disturbance of security motivation. *Psychol. Rev.* **111**, 111-127.
- Szechtman H and EZ Woody (2006) Obsessive-compulsive disorder as a disturbance of security motivation: constraints on comorbidity. *Neurotox. Res.* **10**(2), 103-112.
- Taylor S, D McKay and JS Abramowitz (2005) Is obsessive-compulsive disorder as a disturbance of security motivation? *Psychol. Rev.* **112**, 650-657.
- Teichner G, MD Horner and RT Harvey (2001) Neuropsychological predictors of the attainment of treatment objectives in substance abuse patients. *Int. J. Neurosci.* **106**, 253-263.
- Teichner G, MD Horner, JC Roitzsh, J Herron and A Thevos (2002) Substance abuse treatment outcomes for cognitively impaired and intact outpatients. *Addict. Behav.* **27**, 751-763.
- Tollefson GD (1996) Cognitive function in schizophrenic patients. *J. Clin. Psychiatry* **57**(Suppl. 11), 31-39.
- Totterdell S (2006) The anatomy of synaptic interactions in cortical and limbic regions. *Neurotox. Res.* **10**(2), 65-86.
- Totterdell S and GE Meredith (1997) Topographical organisation of projections from the entorhinal cortex to the striatum of the rat. *Neuroscience* **78**, 715-729.
- Totterdell S and AD Smith (1986) Cholecystokinin-immunoreactive boutons in synaptic contact with hippocampal pyramidal neurons that project to the nucleus accumbens. *Neuroscience* **19**, 181-192.
- Totterdell S and AD Smith (1989) Convergence of hippocampal and dopaminergic input onto identified neurons in the nucleus accumbens of the rat. *J. Chem. Neuroanat.* **2**, 285-298.
- Tukel R, V Keser, N Karali, T Olgun and C Calikusu (2001) Comparison of clinical characteristics in trichotillomania and obsessive-compulsive disorder. *J. Anxiety Disord.* **15**, 433-441.
- Tukel R, A Polat, O Ozdemir, D Aksut and N Turksoy (2002) Comorbid conditions in obsessive-compulsive disorder. *Compr. Psychiatry* **43**, 204-209.
- Valenti O and AA Grace (2005) Chronic stress affects medial prefrontal cortex neuron activity. *Neuropharmacology* **49** (Suppl.), 1-28.
- van Minnen A, KA Hoogduin, GP Keijsers, I Hellenbrand and GJ Hendriks (2003) Treatment of trichotillomania with behavioral therapy or fluoxetine: a randomized, waiting-list controlled study. *Arch. Gen. Psychiatry* **60**, 517-522.
- Walker EF and D Difrio (1997) Schizophrenia: a neural diathesis-stress model. *Psychol. Rev.* **104**, 667-685.

- Waugh M, M Jackson, GA Fox, SH Hawke and RR Tuck (1989) Effect of social drinking on neuropsychological performance. *Br. J. Addict.* **84**, 659-667.
- Weinberger DR (1987) Implications of normal brain development for the pathogenesis of schizophrenia. *Arch. Gen. Psychiatry* **44**, 660-669.
- Weinberger DR, KF Berman and RF Zec (1986) Physiological dysfunction of dorsolateral prefrontal cortex in schizophrenia: I. Regional blood flow evidence. *Arch. Gen. Psychiatry* **43**, 114-125.
- Weinberger DR, KF Berman and BP Illowsky (1988) Physiological dysfunction of dorsolateral prefrontal cortex in schizophrenia: III. A new cohort and evidence for a monoaminergic mechanism. *Arch. Gen. Psychiatry* **45**, 609-615.
- Westergaard T, PB Mortensen, CB Pedersen, J Wohlfart and M Melbye (1999) Exposure to prenatal and childhood infections and the risk of schizophrenia: suggestions from a study of sibship characteristics and influenza prevalence. *Arch. Gen. Psychiatry* **56**, 993-998.
- Wilder KE, DR Weinberger and TE Goldberg (1998) Operant conditioning and orbitofrontal cortex in schizophrenic patients: unexpected evidence for intact functioning. *Schizophr. Res.* **30**, 169-174.
- Williams NM, A Preece, G Spurlock, N Norton, HJ Williams, S Zammit, MC O'Donovan and MJ Owen (2003) Support genetic variation in neureglin 1 and susceptibility to schizophrenia. *Mol. Psychiatry* **8**, 485-487.
- Woo TU, RE Whitehead, DS Melchitzky and DA Lewis (1998) A subclass of prefrontal gamma-aminobutyric acid axon terminals are selectively altered in schizophrenia. *Proc. Natl. Acad. Sci. USA* **95**, 5341-5346.
- Woody EZ, V Lewis, L Snider, H Grant, M Kamath and H Szechtman (2005) Induction of compulsive-like washing by blocking the feeling of knowing: an experimental test of the security-motivation hypothesis of obsessive-compulsive disorder. *Behav. Brain Funct.* **1**, 11.
- Woody EZ and H Szechtman (2005) Motivation, time course, and heterogeneity in obsessive-compulsive disorder: response to Taylor, McKay and Abramowitz (2005). *Psychol. Rev.* **112**, 658-661.
- Wright CI and HJ Groenewegen (1995) Patterns of convergence and segregation in the medial nucleus accumbens of the rat: relationships of prefrontal cortical, midline thalamic, and basal amygdaloid afferents. *J. Comp. Neurol.* **361**, 383-403.
- Wright CI, AV Beijer and HJ Groenewegen (1996) Basal amygdaloid complex afferents to the rat nucleus accumbens are compartmentally organised. *J. Neurosci.* **16**, 1877-1893.
- Xiao BG and H Link (1999) Is there a balance between microglia and astrocytes in regulating Th1/Th2-cell responses and neuropathologies? *Immunol. Today* **20**, 477-479.
- Ziedonis D, JM Williams and D Smelson (2003) Serious mental illness and tobacco addiction: a model program to address this common but neglected issue. *Am. J. Med. Sci.* **326**, 223-230.

